


NAME \_\_\_\_\_



Cambridge, 1953. Shortly before discovering the structure of DNA, Watson and Crick, depressed by their lack of progress, visit the local pub.

Problem 1 12 pts \_\_\_\_\_

Problem 9 30 pts \_\_\_\_\_

Problem 2 15 pts \_\_\_\_\_

Problem 10 24 pts \_\_\_\_\_

Problem 3 10 pts \_\_\_\_\_

Problem 11 10 pts \_\_\_\_\_

Problem 4 15 pts \_\_\_\_\_

Problem 12 8 pts \_\_\_\_\_

Problem 5 10 pts \_\_\_\_\_

Problem 13 8 pts \_\_\_\_\_

Problem 6 20 pts \_\_\_\_\_





Problem 14 15 pts \_\_\_\_\_

Problem 7 12 pts \_\_\_\_\_




TOTAL 100 pts \_\_\_\_\_

Problem 8 12 pts \_\_\_\_\_


1. For each pair of molecules below, circle the one that would be the STRONGEST base. (12 pts)








2. Circle all of the following compounds that are AROMATIC. (15 pts)




3. *N*-Methylimidazole is an aromatic heterocycle that contains two nitrogens. Upon treatment with acid, one of these nitrogens is preferentially protonated. On the structure below, draw the protonated form (eg. where does the proton go) and briefly explain why the proton adds selectively to one of the nitrogens. (9 pts)




4. Identify the most acidic hydrogens in each of the following compounds. (15 pts)




5. Briefly explain why the following reaction will not work as shown. What reagents do you need to make the Hell-Vollhard-Zelinskii reaction proceed? (10 pts)




6. Carbonyl condensation reactions can occur between different kinds of carbonyl compounds. The products of three mixed condensation reactions are shown below. Draw the structures of the carbonyl compounds that react to form these products. Make sure to draw the ENOLATE form of the nucleophile and the CARBONYL form of the electrophile. (20 pts)




7. Draw the major product of the following Diels-Alder reactions. (12 pts)




8. Provide the major product of the following reaction sequences. (12 pts)




9. Fill in the missing reagents and structures in the following synthetic sequences. (30 pts)






10. Fill in the missing reagents and structures in the following synthetic sequences. (24 pts)






11. The mechanism for the acid catalyzed hydrolysis of esters shown below has several errors in it. Identify them by circling all incorrect ARROWS and STRUCTURES. (10 pts)







12. Circle the structure that matches the following proton and C13 NMR Data. (8 pts)





13. Circle the structure that matches the following proton and C13 NMR Data. (8 pts)




14. The three compounds shown below have very different  $^{13}\text{C}$  NMR spectra. Match the structures with the correct spectra by placing the letter of the compound in the appropriate box. (15 points)







**Typical NMR Chemical Shifts**  
 (Note - these are approximate and sometimes peaks  
 can show up outside of these ranges).

Functional Group	Type	<sup>1</sup> H Chemical Shift (ppm)	<sup>13</sup> C Chemical Shift (ppm)
$\begin{array}{c}   \\ -\text{C}-\text{H} \\   \end{array}$	Alkane	0.7 - 1.8	10 - 60
$\begin{array}{c}   \\ =\text{C}-\text{C}-\text{H} \\   \end{array}$	Allylic or next to carbonyl	1.6 - 3.0	30 - 60
$\begin{array}{c}   \\ \text{X}-\text{C}-\text{H} \\   \end{array}$	next to halogen or alcohol	2.5 - 4.0	20 - 85
$\begin{array}{c} \text{O} \\    \\ \text{C}-\text{O}-\text{C}-\text{H} \\   \end{array}$	next to oxygen of an ester	4.0 - 5.0	50 - 85
$\begin{array}{c}   \\ =\text{C}-\text{H} \end{array}$	vinyllic	4.5 - 6.5	110 - 150
	aromatic	6.5 - 8.0	110 - 140
$\begin{array}{c} \text{O} \\    \\ -\text{C}-\text{H} \end{array}$	aldehyde	9.7 - 10.0	190 - 220
$\text{O}-\text{H}$	alcohol	varies widely will exchange with D <sub>2</sub> O	N/A
$\begin{array}{c} \text{O} \\    \\ -\text{C}-\text{X} \end{array}$	carbonyl of ester, amide, or carboxylic acid (X = O, N)	N/A	165 - 185
$\begin{array}{c} \text{O} \\    \\ -\text{C}- \end{array}$	carbonyl of ketone or aldehyde	N/A	190 - 220